A Introductory Linear Algebra-I

1. In 3 D a vector has three components:

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k} \tag{A.1}$$

 \hat{i} , \hat{j} and \hat{k} are unit vectors in the x, y and z directions respectively. \hat{i} , \hat{j} and \hat{k} may also be called basis vectors or just bases and this is a terminology that we will use often. Vectors can be represented in the following form:

$$\vec{r} \equiv \begin{pmatrix} x \\ y \\ z \end{pmatrix} \tag{A.2}$$

- 2. The vector is fully described by: (i) its magnitude, and (ii) its direction. Compare this with a scalar number that only has a magnitude.
- 3. The magnitude of the vector \vec{r} can be calculated by performing the following mathematical operation:

$$|\vec{r}| \equiv \sqrt{\vec{r} \cdot \vec{r}} = \sqrt{|x|^2 + |y|^2 + |z|^2}$$
(A.3)

where we have introduced the definition of the "dot" product of two vectors:

$$\vec{a} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}
\vec{b} = b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k}
\vec{a} \cdot \vec{b} = a_1 b_1 + a_2 b_2 + a_3 b_3$$
(A.4)

- 4. The "dot" product is a very fundamental concept, and very useful for our further development hence lets look at it a little closely.
- 5. Note that by the definition of matrix multiplication, I can write the dot product of \vec{a} and \vec{b} as:

$$\vec{a} \cdot \vec{b} = (a_1 \ a_2 \ a_3) \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}$$
(A.5)

To make sure that this is in fact the case, we need to review matrix multiplication. Please do so, or talk to the instructor to make sure that this is not a problem.

6. Homework: Multiply the two matrices:

$$A = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} and B = \begin{pmatrix} 1 & 4 & 2 & 5 \\ 3 & 6 & 9 & 10 \end{pmatrix}$$
(A.6)

Pay careful attention to the order in which these matrices are multiplied. What A multiplied by B? What is B multiplied by A? Based on this exercise what can you say about the order in which matrices are multiplied?

7. Homework: Evaluate Eq. (A.5) using the laws of matrix multiplication. Show that you get the same result as in Eq. (A.4).

A.1 The Dual Space

8. The object $(a_1 a_2 a_3)$ in Eq. (A.5) can be referred to as the *dual-space analogue* of the object:

$$\begin{pmatrix}
a_1 \\
a_2 \\
a_3
\end{pmatrix}$$
(A.7)

- 9. Note that every vector has its *dual-space analogue*, and for real numbers, the dual space analogue is just the *transpose* of the vector.
- 10. So we may write:

$$\vec{a} \cdot \vec{b} = \vec{a}^T \vec{b} \tag{A.8}$$

11. However, in general the components of the vector \vec{a} need not be real (they could be complex!!) and then the dual space analogue is not simply the transpose. In the complex case the dual space analogue of the vector \vec{a} is defined as:

$$\vec{a}^{\dagger} = (a_1^* \ a_2^* \ a_3^*) \tag{A.9}$$

where the "superscripted *" imply *complex conjugation*.

- 12. At this point it is necessary to go ahead and do the homework problems on complex variables before you proceed further with the rest of this handout.
- 13. Homework: Show that the definition in Eq. (A.9) is consistent with the dot product definition in Eq. (A.4). Hint: Evaluate the quantity $\vec{a}^{\dagger}\vec{a}$ using multiplication of matrices as you did in Eq. (A.5). Show that you get the same result as Eq. (A.4).
- 14. The dual space analogue is a very important concept in quantum theory, as we will see later. For now we understand the dual space analogue is the complex conjugated transpose of a vector.
- 15. The basis vectors, \hat{i} , \hat{j} and \hat{k} that we introduced earlier, have a very important property and that is the "dot" product of each of these with itself is equal to 1, but the dot product of one with any of the others is always 0 (zero). This means:

$$\hat{i} \cdot \hat{j} = \hat{i} \cdot \hat{k} = \hat{j} \cdot \hat{k} = 0 \tag{A.10}$$

but

$$\hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k} = 1$$
(A.11)

These relations make the vectors \hat{i} , \hat{j} and \hat{k} an *ortho-normal* basis set. The term *ortho-normal* has two parts: the first part *ortho* is part of the word *orthogonal*. Two vectors are said to be *orthogonal* when they are at 90 degrees. (Note that the vectors \hat{i} , \hat{j} and \hat{k} are at 90 degrees to each other, as is required from the fact that these are unit vectors along the x, y and z directions.) The second part of *ortho-normal* is the word *normal*: a vector is normalized when its magnitude is 1.

B Introductory Linear Algebra-II: *Dyads*

- 1. We shall now introduce a very important mathematical quantity called a *dyad*. This is fundamental in the theory of linear vector spaces, and turns out to be of great importance in quantum mechanics. (Make sure you got the exercise on matrix multplication in the previous section before you proceed into this one.)
 - Before we introduce the dyad, let us reconsider the dot product in Eq. (A.5). The product of a (1×3) matrix with a (3×1) matrix leads to a (1×1) matrix, *i.e.* a scalar number.
 - What would happen if we had things backwards?

$$\begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} (a_1 \ a_2 \ a_3)$$
 (B.1)

This represents a (3×1) matrix multiplied with a (1×3) matrix and hence the result should be a (3×3) matrix !!

- The quantity in Eq. (B.1) is called the *dyadic product* or *outer product* of the two vectors. By contrast the dot product in Eq. (A.5) is also known as the *inner product* of two vectors. (Note, the outer or dyadic product is *not* the cross product, the result of which is a vector, *not* a matrix.)
- 2. To understand some of the implications of the dyadic product (there will be other implications that we will realize when we introduce the Dirac notation) let us consider the basis vectors in 3-dimensional space:

$$\vec{i} = \begin{pmatrix} 1\\0\\0 \end{pmatrix} ; \ \vec{j} = \begin{pmatrix} 0\\1\\0 \end{pmatrix} ; \ \vec{k} = \begin{pmatrix} 0\\0\\1 \end{pmatrix}$$
(B.2)

3. The dyadic product:

$$\vec{i}\vec{i}^{\dagger} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$
(B.3)

that is a matrix. Similarly the dyadic products $\vec{i}\vec{i}^{\dagger}$, $\vec{j}\vec{j}^{\dagger}$, $\vec{j}\vec{i}^{\dagger}$, $\vec{j}\vec{j}^{\dagger}$, $\vec{j}\vec{k}^{\dagger}$, $\vec{k}\vec{i}^{\dagger}$, $\vec{k}\vec{j}^{\dagger}$, $\vec{k}\vec{k}^{\dagger}$ are all matrices with 1 at one of the positions in the matrix and zeroes everywhere else.

4. Homework:

(a) Write down all the dyadic products $\vec{i}\vec{i}^{\dagger}, \vec{i}\vec{j}^{\dagger}, \vec{i}\vec{k}^{\dagger}, \vec{j}\vec{i}^{\dagger}, \vec{j}\vec{j}^{\dagger}, \vec{j}\vec{k}^{\dagger}, \vec{k}\vec{i}^{\dagger}, \vec{k}\vec{i}^{\dagger}, \vec{k}\vec{i}^{\dagger}$

(b) Homework: Show that:

$$\vec{i}\vec{i}^{\dagger} + \vec{j}\vec{j}^{\dagger} + \vec{k}\vec{k}^{\dagger} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$
(B.4)

The matrix on the right hand side in this equation is called the identity or unit matrix. This relation is called the *resolution of the identity* and we will have a great deal of need for this expression.

(c) Using the results obtained above, show that the matrix:

$$\left(\begin{array}{r}
1 & 2 & 3\\
4 & 5 & 6\\
7 & 8 & 9
\end{array}\right)$$
(B.5)

can we written as a linear combination $\vec{i}\vec{i}^{\dagger}, \vec{i}\vec{j}^{\dagger}, \vec{i}\vec{k}^{\dagger}, \vec{j}\vec{i}^{\dagger}, \vec{j}\vec{j}^{\dagger}, \vec{k}\vec{i}^{\dagger}, \vec{k}\vec{i}^{\dagger}, \vec{k}\vec{i}^{\dagger}, \vec{k}\vec{i}^{\dagger}$

- 5. Note: You have just performed a very important exercise. An important essence of quantum mechanics lies in this exercise.
- 6. Firstly, the resolution of the identity is also known as the completeness relation. Any vector can be written as a linear combination of an *ortho-normal* and *complete* set of vectors. As the set \hat{i} , \hat{j} and \hat{k} are *ortho-normal* (as seen in point 15 of Section A) and *complete* (as seen in point 4b) any vector in three-dimension can be written as a linear combination of these vectors.
- 7. Secondly, we also note from point 4 that any matrix is writable as a linear combination of the *dyadic*-basis: $\vec{i}i^{\dagger}$, $\vec{i}j^{\dagger}$, $\vec{i}k^{\dagger}$, $\vec{j}j^{\dagger}$, $\vec{j}k^{\dagger}$, $\vec{k}i^{\dagger}$, $\vec{k}i^{\dagger}$, $\vec{k}k^{\dagger}$.